

Resummations of free energy at high temperature

G. Cvetič*

Department of Physics, Universidad Técnica Federico Santa María, Valparaíso, Chile

R. Kögerler†

Department of Physics, Universität Bielefeld, 33501 Bielefeld, Germany

(Dated: February 1, 2008)

We discuss resummation strategies for free energy in quantum field theories at nonzero temperatures T . We point out that resummations should be performed for the short- and long-distance parts separately in order to avoid spurious interference effects and double-counting. We then discuss and perform Padé resummations of these two parts for QCD at high T . The resummed results are almost invariant under variation of the renormalization and factorization scales. We perform the analysis also in the case of the massless scalar ϕ^4 theory.

PACS numbers: 12.38.Cy, 11.10.Wx, 12.38.Bx, 12.38.Mh

I. INTRODUCTION

During the last decade the physics of many-particle systems at relativistic energies has attracted increasing interest. On the one hand, this was due to the hope that high-energy ion collisions at present day colliders would reach the kinematical regime of the quark-gluon-plasma phase of QCD and would thus give new insight into the phase structure of QCD and into the confinement phenomena. On the other hand, a better understanding of such systems can be applied to ultrarelativistic astrophysical and/or cosmological configurations and should therefore provide a conceptual tool for describing the physics of the early universe, in particular phenomena such as baryon number generation, big-bang nucleosynthesis, etc. Although it is not yet clear to what extent the various mentioned configurations can be considered as being in thermal equilibrium, the first attempts at a systematic analysis concentrated on equilibrium statistical mechanics.

Since we are convinced that the interaction of particles at energies up to 1 TeV is described by the standard model of strong and electroweak interactions, the main task consists in formulating a consistent thermal quantum field theory of QCD and QED. Because QCD is asymptotically free, the corresponding running coupling parameter becomes small at a sufficiently high temperature T ($\alpha_s(T) \ll 1$)¹ and a perturbative treatment seems possible. This expectation is supported by the observation that lattice gauge theory calculations [1, 2, 3] indicate that the physics of the (quark-)gluon plasma at T larger than four times the critical temperature T_c is to a very good approximation that of an ideal gas. In particular, the free energy F of such a system is off the ideal gas value by less than 20%. Even more, if masses with an appropriate temperature dependence are accepted, the lattice results [4] agree almost exactly with the ideal gas ones for all values of $T > T_c$. This would suggest that perturbative treatments remain candidates for a description of the relevant phenomena.

Unfortunately, a straightforward perturbative treatment has led to results which are far from satisfactory. The problems emerging can be seen best in the expressions for the free energy F as calculated in various theoretical models. The most exhaustive calculations, up to $\sim g^5$, have been performed for the massless ϕ^4 theory [5, 6] and for QCD (without and with quarks) [7, 8]. These results have the following unpleasant features:

1. There is a very bad convergence behavior of the series,² some of the higher-order terms in the perturbation series are larger than the leading term unless the coupling parameter is very small – in QCD, the coupling $\alpha_s(T)$ would have to be at $T > 1$ TeV. If successive orders are included in the truncated perturbation series (TPS), the sum takes on alternating large values (cf. Fig. 1 of Ref. [9]).
2. At every fixed order the TPS's show a strong dependence of the (arbitrary) renormalization scale (RScl) μ . This results in an additional severe uncertainty of the evaluation, this time due to the ambiguity of the choice of the

*Electronic address: gorazd.cvetic@fis.utfsm.cl

†Electronic address: koeg@physik.uni-bielefeld.de

¹ T being considered as a measure of the average energy of the system's constituents.

² Formally, the series is believed to be an asymptotically divergent series.

RScl μ . In an asymptotically free theory one usually is on the safe side if one chooses μ close to the (lowest) physically relevant momentum scale typical of the contributing configurations, since otherwise the expansion coefficients would blow up. In the case of a system at temperature T one would be led to choose $\mu \sim T$, or more specifically $\mu = 2\pi T$, i.e., the energy of the first nonvanishing Matsubara mode. In fact, this is the choice actually used in the literature. However, we should keep in mind that in the case of a plasma consisting of massless components there are important physical effects connected with much lower energies (screening of the chromoelectric or chromomagnetic field). Therefore, the choice $\mu \approx 2\pi T$ seems not to be well founded, *a priori*.

These features are related to the rather large genuine collective effects. And, in fact, it has been noted recently that these unexpectedly large corrections are rooted in the terms $\sim g^3, g^5$ within the expansions. Such terms render the expression nonanalytic in $a \equiv g^2/(4\pi^2)$ and they do not emerge in the ordinary ($T=0$) field theory. They are the result of a certain (partial) resummation which is necessary to get rid of these infrared divergences [11, 12]. They are closely related with collective effects such as screening or Landau damping.

This observation has constituted the basis for several attempts to remedy the situation: since a specific method of resummation seems to be the root of the problems, one is led to try different resummation procedures. The main attempt goes under the name of screened perturbation theory (SPT) [10, 12] and has been made more systematic by what is called optimized perturbation theory [13]. The main idea is to add a local mass term to the free part of the Lagrangian and subtract it from the interaction part. The former is treated nonperturbatively, constituting a genuine (screening) mass; the latter perturbatively. Physically this means that one expands about a (ideal) gas of massive quasiparticles (dressed gluons, for example) rather than about massless particles. The technical consequences are striking. Since the objects one starts with are massive, there is no infrared problem and no need for resummation, and the resulting expressions have better convergence behavior than the original perturbation series. However, for obtaining numerical results, one has to fix the chosen mass at the final stage. The authors of Ref. [12], e.g., used (an appropriate approximation of) the gap equation for fixing this mass.

All this is relatively straightforward in a theory such as ϕ^4 where nothing forbids a genuine mass term. Within a gauge theory like QCD the whole procedure becomes much more cumbersome, since the addition of a genuine (local) mass term would spoil the gauge symmetry from the outset. A technical way out of this problem is hard-thermal loop (HTL) perturbation theory [9]. This is a SPT generalization which respects gauge invariance. The procedure again rests on a (partial) resummation: Those higher-order loop corrections are included that are of leading order in g_s for amplitudes involving soft external momenta $p \sim g_s T$. Unfortunately, the resulting HTL correction terms are nonlocal. This leads to complicated UV divergences, and only some of them are canceled by physical mechanisms (quasiparticle formation and Landau damping). The rest have to be tamed by artificial counterterms proportional to the quasiparticle mass and they generate an additional renormalization scale dependence. Despite this complicated and not uniquely specified procedure, the resulting series has again a better convergence behavior than the original perturbation series. On the other hand, when the so called Φ -derivable approximation scheme [14, 15] is applied, some of the problems of the HTL perturbation theory are avoided. In this scheme, the HTL contributions to the free energy were resummed in Ref. [15] in ϕ^4 theory, QED, and QCD, but the UV divergences were shown to cancel. A related approach which uses the framework of the scalar $O(N)$ -symmetric model in the large- N limit was developed in Ref. [16]. Another approach, overcoming the infrared problems appearing in the conventional perturbation theory and using dressed propagators, was developed in Ref. [17].

On the other hand, the strong renormalization scale (RScl) dependence is not or not sufficiently reduced by the SPT or HTL approach. Further, there remains a principal question: Why are the SPT results so much better than those of the naive perturbation approach? Although the need for resummation is avoided in the first step in SPT due to the mass m_g of the quasiparticle, the correct choice of the specific formula for m_g again needs a resummation implicitly contained in the gap equation. As long as these points are not clarified, the SPT treatment cannot be considered fully satisfying.

A completely different way of remedying the weak points of the naive perturbation theory consists in replacing the truncated perturbation series (TPS) by appropriate Padé approximants (PA's). This approach is motivated by at least two features of PA's: First, PA's at increasing order in general show much better convergence than the TPS's from which they are obtained. It has been shown that even when the TPS's are divergent (asymptotic series), the corresponding PA's may converge and do it under rather general conditions [18]. Secondly, PA's reduce considerably the RScl dependence of the TPS. In fact, it is known that the diagonal PA's, constructed from a TPS of an RScl-independent quantity in powers of $a(\mu) \equiv g^2(\mu)/(4\pi^2)$, are RScl-independent in the limit of the one-loop running of the coupling parameter $a(\mu)$ (large β_0 -limit) [19]. Further, related approximants have been developed which are exactly RScl-independent [20] and even renormalization scheme independent [21]. Since the full perturbation series corresponds to a physical (in principle measurable) quantity, this quantity is exactly RScl-independent. Therefore, one is led to conjecture that PA's and related approximants are nearer to the true value than the original TPS's. The physical reason for the (approximate) RScl independence of such approximants is that they include a certain resummation [19], thus containing infinitely many terms whose absence was responsible for the

spurious RScl-dependence of the TPS.

The first applications of PA's to thermal perturbation theory were made by Kastening [22] and by Hatsuda [23]. These authors started from the available TPS for the free energy F (in ϕ^4 and QCD), which is a TPS in powers of $a^{1/2} = g/(2\pi)$, up to $\sim a^{5/2}$, and replaced it by various PA's based on it. They demonstrated improved RScl stability of these PA's. Although the relatively large number of powers in g of the underlying TPS suggests a good convergence quality of the resulting PA's, the results have to be treated with caution. First, there is no formal reason to expect that certain PA's, specifically the diagonal ones, should be stable under variation of the RScl. In this respect, we note that it is the one-loop (large- β_0) running of $a = g^2/(4\pi^2)$,

$$a(\mu_1) = \frac{a(\mu_0)}{1 + a(\mu_0)\beta_0 \ln(\mu_1^2/\mu_0^2)} , \quad (1)$$

which is responsible for the (large- β_0) RScl independence in diagonal PA's of underlying TPS's in powers of a . This is so because relation (1) represents a homographic transformation $a \mapsto a/(1 + Ka)$ [19]. However, relation (1) yields a structurally different one-loop running of $g = 2\pi a^{1/2}$, namely

$$g(\mu_1) = \frac{g(\mu_0)}{[1 + g^2(\mu_0)\beta'_0 \ln(\mu_1^2/\mu_0^2)]^{1/2}} , \quad (\beta'_0 \equiv \frac{\beta_0}{4\pi^2}) . \quad (2)$$

This relation is not homographic, and consequently the PA's in such a case need not be (large- β_0) RScl-invariant.

There is a second, more significant point of criticism of the aforementioned PA approach. It can be best explained in terms of diagrams. As we have mentioned, the PA's represent a resummation (analytic continuation) of the infinite sum of a certain class of diagrams. On the other hand, the TPS expressions for $F(T)$ also include a selective resummation (of ring and super-ring diagrams). Therefore, a naive application of PA's to the original TPS constitutes a mixing of (at least) two inequivalent resummation effects and could easily lead to partial double-counting or other (interference-like) inconsistencies. In the context of QED and QCD at $T = 0$, somewhat similar aspects have been pointed out and accounted for in Refs. [24].

Within the present paper we want to stick to PA's because of their unique advantages: reduced RScl-dependence, better convergence properties, and (quasi)analytic continuation aspects. However, we are going to develop a procedure that is free of the aforementioned weaknesses. We do this by separating all the terms in the full available TPS into groups which represent TPS's of separate physical, i.e., RScl-independent, quantities. To each of the TPS's we apply PA's separately. The resulting expressions are not only (approximately) RScl-independent, but are supposed to represent better approximations to the true values since double-counting and interference-like inconsistencies are excluded.

In Sec. II we present the main elements of our method. In Sec. III we apply this method to QCD, and in Sec. IV to the massless ϕ^4 theory. Section V summarizes the results and presents conclusions.

II. SEPARATION OF LONG AND SHORT DISTANCE REGIMES

We start with the perturbatively calculated expressions for the free energy density F both for massless ϕ^4 and for QCD. They have been calculated recently [5, 6, 7, 8] up to $\sim g^5$, i.e., three loops plus all the ring diagrams summed up. The generic structure of the resulting expressions is

$$F(T) = F_{\text{ideal}} \left[C_0 + C_2 a + C_3 a^{3/2} + C_4(\mu) a^2 + C_5(\mu) a^{5/2} + \mathcal{O}(a^3 \ln a) \right] . \quad (3)$$

Here, a is short-hand for $a(\mu) = g^2(\mu)/(4\pi^2)$, i.e., the running (RScl-dependent) coupling parameter. The renormalization scheme (RSch) is assumed to be fixed, say $\overline{\text{MS}}$. The RScl-running is described by the perturbatively specified renormalization group equation (RGE)

$$\frac{\partial a(\mu^2)}{\partial \ln \mu^2} = -\beta_0 a^2 - \beta_1 a^3 - \beta_3 a^4 - \dots . \quad (4)$$

It is interesting to note that in the series (3) the coefficients C_0, C_2, C_3 do not depend on the RScl μ , but C_4 and C_5 do, the μ -dependence showing up as an additive contribution $\propto \ln[\mu/(4\pi T)]$ to the corresponding coefficient. In QCD, C_4 includes a term $\propto \ln[a(\mu)]$ [$\sim 1/\ln \ln(\mu)$]; in ϕ^4 theory, C_5 includes a term $\propto \ln[a(\mu)]$. It is crucial to keep in mind the origin of the various terms: The terms $\sim a^{3/2}$ and $\sim a^{5/2}$ come exclusively from the resummation of the ring diagrams that is necessary to avoid infrared divergences [5, 7]. This resummation procedure also yields part of the

coefficient C_4 of the $\sim a^2$ term. In particular, the contributions $\propto \ln(a)$ in C_4 and C_5 are generated by this procedure as the remnant of the emerging dependence on the screening mass. The other terms (C_0, C_2 , and the remaining part of C_4) stem from ordinary perturbation theory in powers of a and do not contain resummation effects. We further note that the term $\sim a^3$ ($\sim g^6$) cannot be obtained by perturbative methods (which include ring summation) because of the severe infrared divergences appearing at that order [25].

Our aim is to apply Padé approximants (PA's) in a consistent manner to a TPS of the type (3). As argued in the Introduction, in order to avoid an uncontrollable mixing and superposition of different resummations, the separation of the pure perturbative from the ring resummation-generated terms should be performed. In addition, however, we want to take advantage of the approximate μ -independence of the PA's when they are applied to TPS's (in a) of μ -independent quantities. Therefore, the right-hand side of Eq. (3) should be split so that the resulting parts (infinite power series) represent quantities that are μ -independent ("physical") separately. This suggests that a physical principle of separation should be involved, i.e., one which is connected with measurable effects.

In finding such a separation principle one is guided by the decomposition of all thermodynamic quantities in (Fourier) modes. Within the imaginary time formalism, the free energy F at every given order is expressed as a sum over the Matsubara frequencies ω_n

$$\omega_n = 2\pi T n \quad \text{for bosons ,} \quad (5)$$

$$= \pi T(2n + 1) \quad \text{for fermions ,} \quad (6)$$

where $n = 0, 1, \dots$. Since the thermodynamic quantities and correlation functions can be derived from F , they also show up as sums over modes. For a given correlation function the contribution from the (exchange of the) Fourier mode with frequency ω_n falls off at large spatial distances as $\exp(-\omega_n R)$ (at least if T is larger than all contributing masses). Therefore, the only mode which does not produce an exponentially vanishing contribution to the long-range correlation functions is the bosonic zero mode $\omega_0 = 0$ [Eq. (5)]. Consequently, at sufficiently high T , the static correlators of the contributing fields at large distances $R \gg 1/T$ are exclusively determined by the zero mode. Since this long-distance behavior of correlators is, at least in principle, observable ("physical"), the procedure of separating the bosonic zero-mode (long-distance) contributions from all the other (short-distance) ones rests on physical grounds. We therefore expect that both contributions are separately μ -independent since both have a physical meaning. Further, we know that all resummation effects in the series (3) contribute exclusively to the long-distance part. This is so because these resummation effects of the ring diagrams exclusively stem from the zero-mode contributions, which represent the strongest infrared divergences at the single diagram level. The long-distance part, i.e., the resummed ring diagrams, shows up as a power series in powers of $g \propto a^{1/2}$, starting with the $a^{3/2}$ term. The short-distance part, on the other hand, has the ordinary perturbation character – a perturbation series in integer powers of $a \equiv g^2/(4\pi^2)$.

The discussed decomposition of F represents the basis for our improvement of the underlying TPS results: We apply the appropriate PA's to the separate parts, so that (at least some of) the approximants are approximately RScl-independent, and presumably better converging. In the case of the short-range contributions these are the diagonal PA's $[n/n](a)$. At the available order of the underlying TPS (3), the short-range part is of the form $F_S(T) = F_{\text{ideal}}(C_0 + C_2 a + \bar{C}_4 a^2)$, and the only possible diagonal PA is $[1/1](a)$.³ In the case of the long-range contributions, one tries to see which PA $[n/m](g)$ ($n + m = 2$) is approximately RScl-independent.

Before continuing, we mention an alternative way to interpret the described decomposition on physical grounds: The long-distance contribution reflects the observable phenomena of screening of interactions and formation of quasi-particles (collective modes). To understand this, we recall the following facts. It is with the resummation of the ring diagrams that the zero modes acquire a (screening) mass $m_s \sim gT$ (m_s can be defined as the solution of the gap equation [10, 12]). And vice-versa: It is the screening phenomenon which explains the striking fact that, while all individual ring diagrams are infrared divergent, their total sum is convergent. From this one concludes that, due to the observable nature of the screening phenomenon, the long-range part F_L , which encompasses the resummation-generated effects, is of physical nature and should therefore be RScl-independent. This can best be seen by determining the screening mass via the gap equation [10, 12] which is equivalent to summing up the ring diagrams.

This interpretation also shows that in theories with more degrees of freedom (like QCD) there might be several stages of screening, e.g., screening of chromoelectric and chromomagnetic gluons, respectively. Consequently, the free energy F can then be decomposed into more than two parts: the short-distance part, and one part for each kind of screening.

Having described the physical idea behind our approach, we have not yet addressed the technical problem of actual decomposition. In principle, the answer is simple: in order to single out the long-range (zero-mode) contribution, one

³ The Padé approximant $[n/m](x)$ for a quantity $S(x)$ is a ratio of two polynomials of n 'th and m 'th degree in x : $[n/m](x) = P_n(x)/P_m(x)$, such that its expansion in powers of x reproduces the terms up to $\sim x^{m+n}$ of the expansion of $S(x)$: $S(x) = r_0 + r_1 x + \dots r_{n+m} x^{n+m} + \dots$

has to integrate out all higher modes (bosonic, and all fermionic modes). The resulting expression, when subtracted from the full expression, should yield the short-distance contribution, i.e., that of all the nonzero modes. In practice, integrating out explicitly all higher modes is a cumbersome task. Fortunately, there exists an alternative method, the method of effective field theories, which was developed for thermal perturbation theory by Braaten and Nieto [6, 8, 26]. The method can be briefly described as follows. At small distances ($R \leq 1/T$) the behavior of the system is determined by ordinary perturbative QCD. On the other hand, the long-distance behavior is dominated by the zero modes and can, consequently, be described by an effective field theory which, at large T , is a bosonic field theory in three dimensions ("dimensional reduction" [25, 27]). The effective bosonic field is approximately identified with the zero modes of the original fields. For the construction of the effective field theory one does not need to specify this effective (static) boson field in terms of the original fields exactly. One simply writes down the most general three-dimensional Lagrangian for the effective fields that respects the symmetries of the original theory. In general, this effective Lagrangian contains infinitely many terms – operator expressions of arbitrary high dimensions – and is thus nonrenormalizable. Therefore, an ultraviolet cutoff Λ is needed. The corresponding effective coupling parameters $g_{E,i}$ ($i=1, 2, 3$; E stands for effective) are then determined by a matching procedure: One computes sufficiently many static correlation functions, both in the original and in the effective theory, with as yet unspecified $g_{E,i}$, and demands that the results agree at larger distances $R > 1/T$. The resulting matching relations allow one to express the effective $g_{E,i}$'s in terms of the original coupling parameter g , T , and the cutoff Λ :

$$g_{E,i} = g_{E,i}(g, T, \Lambda) . \quad (7)$$

The cutoff Λ ($\sim T \sim 2\pi T$), also called the factorization scale, is roughly the momentum below which the effective theory should take over. Relations (7) can be understood as perturbation series in the coupling parameters g . Fortunately, the $g_{E,i}$'s corresponding to interaction operators with higher dimensions are of higher order in g . Therefore, for calculation at a given order only a very restricted number of $g_{E,i}$'s has to be taken into account.

The effective theory approach has been applied by Braaten and Nieto to ϕ^4 theory [6] and to QCD [8].

In the case of $g^2\phi^4/4!$ there is only one cutoff Λ separating the long- and short-distance regimes [6]. Consequently, the free energy density F consists of two parts

$$F = F_S + F_L , \quad (8)$$

where the long-range part F_L is determined by the effective theory and contains the effective coupling parameters $g_{E,i}$. If these $g_{E,i}$'s are expanded in powers of g , then Eq. (8) yields the formula for F that was originally obtained perturbatively by including the ring-diagram resummation [5, 7]. It is a check of consistency of the effective field approach to show that the total expression for F is independent of the factorization scale Λ .

In QCD one can separate from the short-distance regime two long-distance regimes (chromoelectric and chromomagnetic) [8] corresponding to two types of screening. Therefore one can apply two effective theories. One is called the electrostatic QCD (EQCD), and contains electrostatic and magnetostatic effective gauge fields. It describes the physics of the quark-gluon system at distances $r \gtrsim 1/m_E$, where $m_E \sim gT$ is the mass scale of the chromomagnetic screening. The other is called magnetostatic QCD (MQCD), and contains only magnetostatic fields. It acts at distances $r \gtrsim 1/m_M$, where $m_M \sim g^2T$ is the magnetic screening mass, which, however, cannot be calculated perturbatively because of severe infrared divergences at the order g^6 [25, 28]. As a consequence, the free energy density F of hot QCD consists of three contributions:

1. the short-distance [$r \lesssim 1/(\pi T)$] contribution F_E of nonzero modes, i.e., the modes with frequencies equal to and higher than the first Matsubara frequency $\omega_1 = 2\pi T$;
2. the long-distance contribution F_M , with $r \sim 1/m_E \sim 1/(gT)$, i.e., the collective modes described by EQCD effective theory; and
3. the "rest" contribution F_R from even larger scales $r \sim 1/m_M \sim 1/(g^2T)$ of MQCD effective theory.

Each of these three contributions is expected to be physical, i.e., μ -independent.

III. THE CASE OF QCD

A. Formulas for the approach

We will first apply our approach to QCD since this is physically and experimentally the most interesting case. The free energy density F , for an arbitrary number n_f of quark generations, has been calculated in thermal perturbation

theory up to terms $\sim a^{5/2}$ [7, 8], within the $\overline{\text{MS}}$ renormalization scheme. As mentioned before, the free energy density can be decomposed into three physically distinct parts [8]

$$F = F_E + F_M + F_R . \quad (9)$$

Up to order $a^{5/2}$, only F_E and F_M contribute, and they are, in principle, perturbatively computable. These are the contributions from the energy intervals (Λ_E, ∞) and (Λ_M, Λ_E) , respectively, where the factorization scales Λ_E and Λ_M delimit the ultraviolet (UV) and infrared (IR) bounds of EQCD and satisfy the inequalities

$$m_M(\sim g_s^2 T) \lesssim \Lambda_M \lesssim m_E(\sim g_s T) \lesssim \Lambda_E \lesssim \omega_1(=2\pi T). \quad (10)$$

Apart from fitting into this hierarchy, the factorization scales Λ_E and Λ_M are in principle arbitrary. The sum of the three contributions is independent of these scales, and $F_E + F_M$ is independent of Λ_E . Once Λ_E and Λ_M are fixed, each of the three contributions is a (quasi)observable, i.e., a quantity that is independent of the renormalization scale and scheme. We will show that the specific RScI-dependence of the available TPS's for F_E , m_E , and F_M is in fact consistent with RScI-independence of the full quantities F_E , m_E , and F_M .

Specifically, we have for F_E the following TPS [8]:

$$F_E = -\frac{8\pi^2}{45} T^4 \left[\left(1 + \frac{21}{32} n_f\right) - \frac{15}{4} \left(1 + \frac{5}{12} n_f\right) \tilde{F}_E(\Lambda_E) \right] , \quad (11)$$

where the first term represents the contribution of the free (ideal) quark-gluon gas

$$F_{\text{ideal}} = -\frac{8\pi^2}{45} T^4 \left(1 + \frac{21}{32} n_f\right) , \quad (12)$$

and the “canonical” QCD-part $\tilde{F}_E(\Lambda_E)$ is

$$\tilde{F}_E(\Lambda_E) = a(\mu^2) \left\{ 1 + \left[Q(n_f; \Lambda_E) + L(n_f) \ln \left(\frac{\mu^2}{4\pi^2 T^2} \right) \right] a(\mu^2) \right\} + \mathcal{O}(a^3) , \quad (13)$$

with

$$Q(n_f; \Lambda_E) = -18 \frac{(1+n_f/6)}{(1+5 n_f/12)} \ln \left(\frac{\Lambda_E^2}{4\pi^2 T^2} \right) + \frac{4}{15} \frac{(-244.898 - 17.2419 n_f + 0.415029 n_f^2)}{(1+5 n_f/12)} , \quad (14)$$

$$L(n_f) = \frac{11}{4} \left(1 - \frac{2 n_f}{33} \right) . \quad (15)$$

Here,⁴ n_f is the number of active quark flavors; $a(\mu^2) \equiv \alpha_s(\mu^2; \overline{\text{MS}})/\pi \equiv g_s^2(\mu^2; \overline{\text{MS}})/(4\pi^2)$. The RScI μ is usually chosen as $\sim \pi T$. We note that $L(n_f)$, the coefficient of the $(\ln \mu^2)$ -dependent part at order a^2 , is exactly equal to β_0 , the one-loop coefficient of the QCD beta function of the RGE (4). It is then exactly this fact which guarantees that the derivative $\partial \tilde{F}_E / \partial \ln \mu^2$ has the terms of $\sim a^2$ canceled due to the RGE. Therefore, this variation is $\sim a^3$, which is of the order of the first unknown term in the TPS of $\tilde{F}_E$. This shows that the TPS behavior of $\tilde{F}_E$ (and F_E) is compatible with the supposition of the μ -independence of F_E .

Further, the available TPS for the electric Debye-screening mass m_E is [8]

$$\tilde{m}_E^2 \equiv \frac{1}{4\pi^2 T^2} \frac{1}{(1+n_f/6)} m_E^2 = a(\mu^2) \left\{ 1 + \left[P(n_f) + \beta_0 \ln \left(\frac{\mu^2}{4\pi^2 T^2} \right) \right] a(\mu^2) \right\} + \mathcal{O}(a^3) , \quad (16)$$

where

$$P(n_f) = (0.612377 - 0.488058 n_f - 0.0427979 n_f^2) / (1 + n_f/6) . \quad (17)$$

Again, it is possible to see that $\partial \tilde{m}_E / \partial \ln \mu^2 \sim a^3$, compatible with the μ -independence of this quantity.

⁴ We note that there are two misprints in a formula for F_E ($f_E \equiv F_E/T$) in Refs. [8] – in Eq. (7) (PRL) and Eq. (54) (PRD) – in the sign of the coefficient at $\ln[\Lambda_E/(2\pi T)]$ and in the sign of the term $17.24 n_f$.

Let us now consider the part F_M , which includes all long-distance contributions, i.e., all ring-diagram resummation effects. The available TPS for F_M was calculated within EQCD (with effective mass m_E and effective coupling g_E) in Ref. [8]. Since $g_E^2(\mu) = g_s^2(\mu)T [1 + \mathcal{O}(g^2)]$, one can express the result immediately in powers of g_s . This gives

$$F_M = \frac{2}{3\pi} T m_E^3 \tilde{F}_M(\Lambda_E; \Lambda_M), \quad (18)$$

with

$$\begin{aligned} \tilde{F}_M(\Lambda_E; \Lambda_M) = & -1 + \left(0.255838 + \frac{9}{4} \ln \frac{\Lambda_E^2}{m_E^2} \right) \frac{1}{2\pi} (1+n_f/6)^{-1/2} g_s(\mu^2) \\ & + 27.5569 \frac{1}{4\pi^2} (1+n_f/6)^{-1} g_s^2(\mu^2) + \mathcal{O}(g^3). \end{aligned} \quad (19)$$

The Λ_M -dependence would first show up at terms of higher order $\sim g_s^3$. The Λ_E -dependence cancels in the sum $F_E + F_M$ to the orders available in Eqs. (11)-(15) and (18)-(19). Equation (19) is a series in $g_s(\mu^2) \equiv 2\pi\sqrt{a(\mu^2)}$. To the order available, it is automatically compatible with μ -independence of F_M since $\partial g_s(\mu^2)/\partial \ln \mu^2 \sim g_s^2$ by the RGE (4).

When the screening mass m_E (16) is expanded in powers of $g_s(\mu)$ and inserted in Eqs. (18)-(19), a power expansion of F_M in $g_s(\mu)$ can be obtained, starting with g_s^3 ($\sim m_E^3$)

$$\begin{aligned} F_M = & \frac{2}{3\pi} k_f^3 T^4 \left\{ -g_s^3(\mu^2) + \frac{1}{2\pi k_f} \left[0.255838 - \frac{9}{2} \ln(g_s(\mu^2) k_f) + \frac{9}{2} \ln(\Lambda_E/T) \right] g_s^4(\mu^2) \right. \\ & \left. + \frac{1}{4\pi^2 k_f^2} \left[-\frac{3}{2} \left(P(n_f) + \beta_0 \ln \left(\frac{\mu^2}{4\pi^2 T^2} \right) \right) k_f^2 + 27.5569 \right] g_s^5(\mu^2) + \mathcal{O}(g_s^6 \ln g_s) \right\}, \end{aligned} \quad (20)$$

where we used the notation $k_f = (1+n_f/6)^{1/2}$ and Eq. (17). Adding it to the expansion (11)-(13) for F_E , a TPS for F_{E+M} in powers of $g_s(\mu)$ up to $\sim g_s^5$ is obtained as given in Ref. [7] (third entry) and in Ref. [8] (second entry), i.e., a TPS of the form (3). Interestingly, due to the use of the expansion of m_E , the coefficient $C_4(\mu)$ becomes g_s -dependent [dependent on $\ln g_s^2(\mu)$], which represents an additional, although possibly only formal, obstacle for the direct application of PA's to such a TPS.⁵ We avoid this problem by using in the free energy the Padé-resummed squared screening mass m_E^2 .

It is well known that the TPS (19) for $\tilde{F}_M$ in QCD has very bad divergent behavior, and this is the case to a somewhat lesser extent for the TPS of m_E . Thus, their direct evaluations do not yield useful predictions. Of course, this bad divergent behavior is transported to F_M and to $F = F_E + F_M$. To remedy this, the authors of Refs. [22, 23] presented evaluations of F via various Padé approximants (PA's), which were based on the TPS of the expansion of F in powers of g_s , i.e., the sum of Eqs. (11)-(13) and (20), and no separation was performed. Although their results showed significantly reduced μ -dependence of the Padé resummed values of F_{E+M} in comparison to the μ -dependence of the TPS of F_{E+M} , we believe that this approach is not well motivated. This is so because it probably leads to partial double-counting or other interference-like inconsistencies, as argued in the previous sections. We will illustrate this argument with the following simple example. Suppose that we want to resum, by a PA, the sum $S \equiv (S_1 + S_2)$ of two observables S_1, S_2 . Suppose that we have these two observables available as power series of a up to next-to-leading order (NLO):

$$S_j = a(1 + r_1^{(j)} a) + \mathcal{O}(a^3) \quad (j=1,2) \quad (21)$$

$$\Rightarrow S = 2a \left[1 + \frac{1}{2}(r_1^{(1)} + r_1^{(2)})a \right] + \mathcal{O}(a^3). \quad (22)$$

Applying to this TPS of the sum a PA, say $[1/1]$, and expanding the resummed result back in powers of the coupling a , we obtain

$$S^{[1/1]} = 2a \left[1 - \frac{1}{2}(r_1^{(1)} + r_1^{(2)})a \right]^{-1} \quad (23)$$

$$= 2a \left[1 + \frac{1}{2}(r_1^{(1)} + r_1^{(2)})a + \frac{1}{4}(r_1^{(1)2} + r_1^{(2)2} + 2r_1^{(1)}r_1^{(2)})a^2 \right] + \mathcal{O}(a^4). \quad (24)$$

⁵ Formally, PA's are constructed for TPS's where the coefficients are independent of the expansion parameter.

The coefficient at a^3 of the power expansion of the result has a term $2r_1^{(1)}r_1^{(2)}$, indicating that the result contains mixing effects at $\sim a^3$. This indicates some kind of interference effect between the two amplitudes for the observables (processes) S_1 and S_2 . This is not acceptable because S is the sum of these two independent observables, i.e., the two contributions should be summed up incoherently. This argument remains basically unchanged when different PA's are applied, and/or when the TPS's are of higher order, either in a or in g_s .

It is true that expansion of F_{E+M} in powers of g_s (using also expansion of m_E in g_s) gives us a TPS of relatively high order ($\sim g_s^5 \sim a^{5/2}$), and that we can thus apply PA's of relatively high order. Further, the higher order PA's are known to possess in general a weaker μ -dependence than the corresponding TPS [19]. However, such an approach would predict nonphysical higher order effects and thus lead to unreliable predictions for the sum (F_{E+M}).

As argued in the previous sections, we follow a more conservative approach, by summing up by PA's each TPS (for F_E , m_E , F_M) separately. This approach gives us an additional freedom – to choose the RScI in each of these TPS's separately, in the natural range $\mu \sim 1/r$ where r is a typical distance associated with each observable. For TPS's of F_E , m_E , F_M this implies the RScI choices: $\mu_E \sim \omega_1 (= 2\pi T)$; $\mu_m \sim m_E$; and $\mu_M \sim m_E$. Although, in this approach, we have to take as bases the TPS's (13), (16), (19) of very low order (NLO), and some of these TPS's show very divergent behavior, we will now show that the results nonetheless show remarkably weak dependence on μ and Λ_E .

B. Numerical results

In our numerical analysis we used, unless otherwise stated, for the number of active (massless) quark flavors $n_f = 3$, and for the QCD coupling constant the reference value $\alpha_s(M_\tau^2, \overline{\text{MS}}) = 0.334$. We used the $\overline{\text{MS}}$ scheme, and for the β -function we used, for definiteness, the $[2/3]$ PA.⁶

In Fig. 1 we present the results for the screening mass m_E as a function of the corresponding RScI μ_m , for the temperature choice $T = 1$ GeV. All approximations are based on the (NLO) TPS (16). We see that the diagonal PA $[1/1](a)$ significantly reduces the RScI-dependence, in comparison to the LO and NLO TPS's, in accordance with the arguments of Ref. [19]. Furthermore, the effective charge (ECH) method [29, 30, 31] of fixing the RScI in the NLO TPS gives us a value (fixed by definition) not far from the PA $[1/1]$ values.⁷ We will choose the “physical” screening mass to be determined by the condition $m_E^{[1/1]}(\mu_m) = \mu_m (\equiv m_E^{(0)})$, which gives us the value $m_E^{(0)} = 1.9$ GeV for $T = 1$ GeV.

In Figs. 2(a) and 2(b), we present the results for F_E and F_M as functions of the respective RScI's μ_E and μ_M . The factorization scale Λ_E was chosen, in the spirit of the hierarchy relations (10), to be the logarithmic mean of the typical scales $\mu_E \sim 2\pi T$ and $\mu_M \sim m_E$: $\Lambda_E = \sqrt{2\pi T m_E^{(0)}}$, where $m_E^{(0)} = 1.9$ GeV is the aforementioned “physical” $[1/1](a)$ screening mass. The PA $[1/1](a)$ for F_E is based on the NLO TPS (11)-(13). We see that the PA $[1/1]$ for F_E has drastically reduced the RScI-dependence, and its values are very close to those of the ECH prediction.

On the other hand, the situation with F_M is less favorable, as seen from Fig. 2 (b). The two PA's $[1/1](g_s)$ and $[0/2](g_s)$ were constructed from the TPS in powers of g_s (not a) of Eq. (19), taking for m_E , appearing in Eqs. (18) and (19), the PA $[1/1](a)$ with the RScI's for m_E and F_M taken equal: $\mu_m = \mu_M$. In the case of the (NLO) TPS result for F_M , we used for m_E the NLO TPS result as well. While the two PA's give reduced RScI-dependence of F_M , their values differ drastically. In the case of the PA summation of F_E and m_E the diagonal PA's $[1/1](a)$ are more physically motivated than the off-diagonal ones $[0/2](a)$, due to the RScI-independence of $[1/1](a(\mu))$ in the large- β_0 approximation [19]. That is why we did not employ the $[0/2](a)$ PA's for F_E and m_E . On the other hand, for F_M we have a TPS in $g_s(\mu) = 2\pi\sqrt{a(\mu)}$ of Eq. (19), and not in $a(\mu)$, and therefore the choice $[1/1](g_s)$ for the right-hand side (RHS) of Eq. (19) is not physically better motivated than the choice $[0/2](g_s)$. This can be seen also from the comparably weak RScI-dependence of both PA approximants in Fig. 2(b). In order to choose between the two, we have to study, in addition, their variation under the variation of the factorization scale Λ_E . This will be discussed just below.

In Figs. 3(a) and 3(b), we present the Λ_E -dependence of the approximants. The RScI's were fixed, in accordance with the hierarchies (10), in the following way: $\mu_E = 2\pi T$, $\mu_M = \mu_m = m_E$, where m_E has the aforementioned “physical” value: $m_E = m_E^{(0)} = 1.9$ GeV, for the case of the PA approximants and 1.4 GeV for the case of the TPS's. Figure 3(a)

⁶ As argued in Refs. [21], this PA gives us a reasonable quasianalytic continuation of the $\overline{\text{MS}}$ β -function for values of $\alpha_s(\mu)$ up to $\alpha_s(\mu) = \pi \times 0.32 \approx 1.0$, i.e., for values where the TPS β gives highly unreliable results.

⁷ A generalization of the method of diagonal PA's has been developed [20, 21], which gives complete RScI-independence [20], or RScI- and RSch-independence [21]. However, we have here TPS's available only at the low NLO order, at which the aforementioned approximants basically reduce to those of the ECH method.

shows that the choice $[1/1](g_s)$ for the TPS of Eq. (19) in F_M leads to strong, and thus unphysical, Λ_E -dependence of F_{E+M} . Thus we have to discard the $[1/1](g_s)$ result of the TPS (19). The choice $[0/2](g_s)$ for this part leads, on the other hand, to a result for F_{E+M} which is remarkably stable under variation of Λ_E in the entire interval $m_E^{(0)} \leq \Lambda_E \leq 2\pi T$.

Therefore, we conclude that the choice $[1/1](a)$ for F_E and for m_E , and the choice $[0/2](g_s)$ for the TPS of Eq. (19), i.e., the curve $F_E[1/1]+F_M[0/2]$ in Fig. 3(a), is the least unreliable among all the curves.

The aforementioned properties of various approximants under changes of the RScI's and of Λ_E remain qualitatively the same when we change the value of the temperature. In Figs. 4(a) and 4(b), we present the results for the approximant $(F_E[1/1]+F_M[0/2])/F_{\text{ideal}}$ as a function of the temperature T and of the coupling $g_s(2\pi T) [=2\pi\sqrt{a(2\pi T)}]$. The middle curve (mid Λ) is for the canonical choice of the scales, $\mu_E=2\pi T$ and $\mu_M=\mu_m=m_E$ where m_E is, for each T , determined in the aforementioned way: $m_E^{[1/1]}(\mu_m)=\mu_m$. The factorization scale is taken to be $\Lambda_E=\sqrt{\mu_E\mu_M}$. The lower curve (low Λ) is for the choice of the scales being all at the lower extreme: $\mu_E=\mu_M=\mu_m=m_E=\Lambda_E$, where m_E is determined in the aforementioned way. The upper curve (high Λ) is for the choice of the scales at the upper extreme: $\mu_E=\mu_M=\mu_m=\Lambda_E=2\pi T$. We notice that the curve of the low Λ choice follows well the curve for the canonical (mid Λ) choice in the entire depicted region of T (of g_s). The other choice of PA's ($[1/1](a)$ for F_E and m_E ; $[1/1](g_s)$ for $\tilde{F}_M$ of Eq. (19)) gives values well outside this range – see Fig. 3(a). The TPS's [for F_E , m_E , and for $\tilde{F}_M$ of Eq. (19)] do give us values in qualitative agreement with the lattice results for a specific choice of scales (mid Λ) [see Fig. 3(a)], but the TPS values change drastically when some of these scales, in particular μ_M , change [see Fig. 2(b)]. In Figs. 4(a) and 4(b), we included, for comparison, the values of the TPS up to $\mathcal{O}(g_s^5)$ in powers of $g_s(2\pi T)$, i.e., the sum of the TPS's (11)-(13) and (20). We note that the values for the TPS up to $\mathcal{O}(g_s^4)$ would be above 1.

As all the curves up to this point have been given for the choice of three active massless quark flavors ($n_f=3$), we present in Figs. 5(a) and 5(b), the results for $n_f=0, 2, 3, 4, 6$ (and for the mid Λ choice of the scales). We see that the curve for $n_f=4$ differs little from that for $n_f=3$ in most of the covered parameter space, while the curve for $n_f=0$ differs significantly. However, in the temperature region of particular interest ($T \sim 0.1\text{-}10$ GeV), the choices $n_f=3$ or 4 are expected to be more realistic.

C. Comparison with other approaches

We will now compare the results presented in Figs. 4 and 5 with those of some other approaches.

As mentioned in Sec. III A, Kastening [22] and Hatsuda [23] applied high order PA's to the full TPS for the sum $R \equiv (F_E + F_M)/F_{\text{ideal}}$, i.e.,⁸ TPS in powers of $g_s(\mu)$ including $\sim g_s^5$. Kastening [22] showed that, in the case of value $g_s(T) \approx 1.1$ [$\alpha_s(T) = 0.1$, $n_f = 6$], the application of PA's $[2/2](g_s)$ and $[2/3](g_s)$ reduced the RScI dependence in comparison with the TPS results. At larger values $g_s(2\pi T) \approx 2$ [$\alpha_s(2\pi T) = 1/3$, $n_f = 3$] there was no significant reduction. On the other hand, Hatsuda [23] applied a kind of modified PA's, by postulating that the expressions in the numerator and the denominator of the PA's have zero coefficients for the term $\sim g_s$. The RScI dependence was then shown to be significantly reduced for many such modified PA's ($[2/3]$, $[3/2]$, $[2/4]$, $[4/2]$; with $n_f = 4$). Further, his curves for the R obtained as a function of $\alpha_s(2\pi T)$ were qualitatively similar to our curves in Figs. 4(b) and 5(b), with $R_{\text{min}} = 0.97\text{-}0.98$. However, the differences between his curves were significant, and there was no clear principle to choose any specific one of them. Furthermore, in our Sec. III A we stressed a more physical point of criticism of this approach.

A reliable comparison with lattice results is hampered by the fact that (QCD-)lattice calculations have reproduced thermodynamic quantities (free energy or pressure, entropy, etc.) only for temperatures between the critical temperature T_c and $4.5T_c$ (i.e., for $T \lesssim 1$ GeV). In this temperature region the resulting values for R are less than 0.87 when $n_f = 0$ [1], significantly lower than 1. If the number of (massless or light) quark flavors n_f is larger than zero, the finite cutoff effects are not quite under control and are estimated [3] to increase the calculated $R(n_f)$ by about 15%, giving at $T = 3.5T_c$ (≈ 0.7 GeV) the values $R \approx 0.90 \pm 0.04$ for $n_f=2$ (cf. Fig. 3 of Ref. [3]). On the other hand, our approach, which is based on improved perturbative considerations, is expected to be reliable only for much higher temperatures $T \gtrsim 10$ GeV where the corresponding values of the effective coupling parameters $g_s(2\pi T)$ are not much higher than 1. At such high temperatures we predict $R \approx 0.975$ or higher, for $n_f \leq 4$ [cf. Fig. 5(a)], i.e., very near to 1. These predictions are not incompatible with the lattice results, however. For example, if we take from Fig. 3 of Ref. [3] the continuum estimate for p/T^4 (with $n_f=2$), which is given as a band-curve for $T \leq 3.5T_c$ (≈ 0.7 GeV),

⁸ We note that in the thermodynamic limit $F = -p$, where p is pressure. Thus, $R = p/p_{\text{SB}}$ where $p_{\text{SB}} \equiv p_{\text{ideal}}$ is the ideal gas pressure.

and extrapolate it smoothly to $T \gtrsim 10$ GeV, we can obtain a fair agreement with our results. Further, we note that for $T \gtrsim 10$ GeV, $g_s(2\pi T)$ can still be rather high [$g_s(2\pi T) = 1.1$ - 1.2] and the numerical evaluations of perturbative TPS for such $g_s(2\pi T)$ differ from our results significantly. For example, the TPS up to $\mathcal{O}(g^5)$, partly visible in Fig. 4, gives us for $g_s(2\pi T) \approx 1.2$ (1.0) [$\Leftrightarrow T \approx 10$ (100) GeV] the value 0.925 (0.952).

One interesting point in connection with lattice data is the dependence of R on the number n_f of light or massless quark flavors. It is well known that the ideal gas expression for F shows an increase for $|F|$ with increasing n_f , Eq. (12). Further, while lattice calculations show that the ratio $R(n_f) \equiv F(n_f)/F(n_f)_{\text{ideal}}$ also increases with increasing n_f at $T \sim T_c$, they indicate an inversion of the n_f -dependence of R at the highest available T values ($T \approx 4T_c$) [3]. This is in accordance with our finding [see Fig. 5(a)] that at $T \gg T_c$ the ratio R decreases with increasing n_f .

A calculation in the Φ -derivable approximation scheme using hard thermal loop (HTL) propagators [15] gives lower minimal values of R ($R_{\min} \approx 0.82$, see Fig. 3 there, where $n_f = 0$), and the minimum is at a much higher value of g_s ($g_s \approx 2.5$) than our curves [$g_s(2\pi T) \approx 1.1$]. However, the Φ -derivable approximation was performed at the leading-loop order, and the expansion of the result R in powers of g_s underestimates the positive $\sim g_s^3$ term by a factor of $1/4$. This indicates that the correction of this effect would push the results for R higher.

IV. RESUMMATION RESULTS IN THE CASE OF ϕ^4 THEORY

We can apply the same methods of resummation in the massless scalar ϕ^4 theory. In this case, we can use the corresponding TPS results from Ref. [6]. The interaction is

$$\mathcal{L}_{\text{int}} = \frac{g^2}{4!} \phi^4, \quad (25)$$

and the β -function in the $\overline{\text{MS}}$ scheme is known to five loops [32]

$$\frac{\partial a}{\partial \ln \mu} = 3a^2 - \frac{17}{3}a^3 + 32.54a^4 - 271.6a^5 + 2848.6a^6 + \mathcal{O}(a^7), \quad (26)$$

where $a(\mu) \equiv g^2(\mu)/(16\pi^2)$. For definiteness, we choose the PA [3/3] for this β -function, in order to simulate better the running in the large- $g(\mu)$ region. The high energy contribution F_S to the free energy F is

$$F_S(T) = -\frac{\pi^2}{90} T^4 \left(1 - \frac{5}{4} \tilde{F}_S(T) \right), \quad (27)$$

$$\tilde{F}_S(T) = a(\mu^2) \left\{ 1 - \left[3 \ln \left(\frac{\mu}{4\pi T} \right) + 9.29324 \right] a(\mu^2) \right\} + \mathcal{O}(a^3). \quad (28)$$

In contrast with the analogous QCD quantity F_E , F_S does not depend on the factorization scale Λ_F (in QCD: Λ_E) at the available order. The TPS for the Debye screening mass m_L is

$$\begin{aligned} \tilde{m}_L^2(\Lambda_F) &\equiv \frac{3}{2\pi^2 T^2} m_L^2(\Lambda_F) \\ &= a(\mu^2) \left\{ 1 + \left[-3 \ln \left(\frac{\mu}{4\pi T} \right) + 4 \ln \left(\frac{\Lambda_F}{4\pi T} \right) + 5.39289 \right] a(\mu^2) \right\} + \mathcal{O}(a^3). \end{aligned} \quad (29)$$

In contrast with the analogous QCD quantity m_E , the mass m_L does depend on the factorization scale. The low energy contribution F_L to F has the following associated TPS:

$$\frac{12\pi}{T} \frac{F_L}{m_L^3(\Lambda_F)} = -1 + \frac{3}{2} \sqrt{\frac{3}{2}} \frac{1}{4\pi} g(\mu^2) + \left[4 \ln \left(\frac{\Lambda_F}{4m_L(\Lambda_F)} \right) + \frac{9}{2} \right] \frac{3}{2} \frac{1}{16\pi^2} g^2(\mu^2) + \mathcal{O}(g^3). \quad (30)$$

The factorization scale dependence now cancels out within F_L , up to the available order ($\sim g^5$). We have the hierarchy (10) as in QCD, with the substitutions $m_E \mapsto m_L$ and $\Lambda_E \mapsto \Lambda_F$. The entire analysis of the QCD case, as described in the previous section, can now be repeated following the same procedures. For the coupling parameter we chose the reference value $g(2\pi \text{ GeV}) = 4$. For the temperature choice $T = 1$ GeV, the results for the screening mass m_L as a function of the corresponding RScI μ_m are presented in Fig. 6, and for F_S and F_L as functions of the corresponding RScI's μ_S and μ_L in Figs. 7(a) and 7(b). The factorization scale Λ_F was determined similarly as in the QCD case: $\Lambda_F = \sqrt{2\pi T m_L^{(0)}}$, where in turn $m_L^{(0)}$ is the “physical” screening mass determined by the condition $m_L^{[1/1]}(\mu_m; \Lambda_F) = \mu_m$ ($\equiv m_L^{(0)}$), which gives us the values $m_L^{(0)} = 0.882$ GeV and $\Lambda_F = 2.354$ GeV (for $T = 1$ GeV).

The PA curves for F_L in Fig. 7(b) were obtained by applying the corresponding PA's $[1/1](g)$ or $[0/2](g)$ to the RHS of Eq. (30) and employing for m_L the PA $[1/1](a)$ at the same RScL ($\mu_m = \mu_L$). The TPS curve was obtained by using the TPS of the RHS of Eq. (30), using for m_L the (NLO) TPS (29). The $[0/2]$ curve in Fig. 7(b) is much less RScL-dependent than $[1/1]$, but not significantly less than the TPS curve.

The dependence of the results on the factorization scale Λ_F is depicted in Fig. 8. We see that the PA choice $[1/1](a)$ for F_S (and m_L) and $[0/2](g)$ for F_L results in weak Λ_F -dependence which is comparable with the TPS result,⁹ while the $[1/1](g)$ choice again gives unacceptably strong Λ_F -dependence.

When the temperature is changing, so is $g(2\pi T)$ according to the RGE (26) (we use $[3/3]$ PA on the RHS). In Fig. 9(a) we present the results for the choice $[1/1](a)$ (for F_S and m_L) and $[0/2](g)$ [for the RHS of Eq. (30)], and for the NLO TPS choice [for F_S , m_L , RHS of Eq. (30)]. The “middle” values (mid Λ) are chosen for the relevant scales $\mu_S = 2\pi T$, $\mu_L = \mu_m = m_L^{[1/1]}(\mu_m; \Lambda_F)$, and $\Lambda_F = \sqrt{\mu_S \mu_L}$. If these scales are changed to low Λ [$\mu_S = \mu_L = \mu_m = \Lambda_F$ with $m_L^{[1/1]}(\mu_m; \Lambda_F) = \mu_m$], or to high Λ ($\mu_S = \mu_L = \mu_m = \Lambda_F = 2\pi T$), the PA results vary with $g(2\pi T)$ as presented in Fig. 9(b).

V. SUMMARY AND CONCLUSIONS

We have presented here a method to improve the predictive power of perturbative thermal field theory. Predictions obtained by ordinary (truncated) perturbation theory (including a resummation to get rid of finite- T infrared divergences) suffer from serious divergence and renormalization scale (RScL) ambiguity problems. Therefore, a careful reorganization of these series is needed, which converts their physical content into expressions with better convergence behavior and greater stability under the variation of the RScL.

Efforts in this direction have been undertaken by various authors during the last five years and considerable improvement has been achieved. However, most of these attempts have improved either the divergence problem or the RScL-ambiguity, but not both.

We suggest an approach that addresses both problems, but concentrating mainly on the unphysical RScL-dependence. It is based on a physically motivated separation of the free energy density F into parts which are separately RScL-independent since each of them has an empirical significance on its own. This can be achieved because the separation principle is determined by observable effects (screening of the different massless degrees of freedom). In this way one simultaneously obtains a clear separation between terms which include (ring-diagram) resummation effects on the one hand, and purely perturbative contributions on the other hand, which are free of any such resummation. This is gratifying since it allows us to consistently apply Padé approximants (PA's) to the available truncated perturbation series (TPS's) of the individual parts, thereby avoiding the danger of double counting and spurious interferences of contributions from different kinematical regimes. We also Padé resum the (observable) screening mass which appears in the long-distance part of the free energy density. The resulting expression for the free energy density has strongly reduced RScL-dependence. In addition, we expect this method to show a good convergence when applied to TPS's of higher order, based on the well-known behavior of PA's [18].

Due to its close connection with physical effects (screening), we consider our approach to be less *ad hoc* and more physically motivated than some of the previous methods.

As a consequence of the aforementioned separation of the free energy into (two) parts, the underlying TPS's for the construction of PA's were of low order. Regarded from a purely numerical point of view, this should alarm us and we should expect significant (nonphysical) instabilities of the resulting low order PA's under variation of the RScL μ , and instabilities for the sum of the two parts under variation of the factorization scale. However, our results both in QCD and in ϕ^4 theory are remarkably stable under both variations. This confirms additionally that the described separation forms a sound basis for the application of PA's. Further, thus resummed values of $R = F/F_{\text{ideal}}$ turn out to be below the value 1 for a relatively wide interval of the coupling parameter: $g_s(2\pi T) \leq 1.87$ [$\Leftrightarrow \alpha_s(2\pi T) \leq 0.278$], i.e., $T \geq 0.4$ GeV, in QCD with $n_f = 3$; see Fig. 4(b). However, the method probably breaks down already at lower $g_s(2\pi T) \approx 1.0$ -1.2, i.e., $T \approx 10$ -100 GeV, where it gives the local minimum $R_{\text{min}} \approx 0.977$ (if $n_f = 3$). The results at $T \gtrsim 10$ GeV do not contradict the results of lattice calculations.

⁹ The latter is Λ_F -dependent because we apply in Eq. (30) in this case the (NLO) TPS value for m_L .

Acknowledgments

The work of G.C. was supported by the FONDECYT (Chile) Grants No. 1010094 and 7010094. R.K. would like to thank the Department of Physics at the UTFSM, Valparaíso, for a very warm hospitality during the course of this work. We are grateful to O. Espinosa and D. Bödeker for helpful discussions.

-
- [1] G. Boyd, J. Engels, F. Karsch, E. Laermann, C. Legeland, M. Lütgemeier, and B. Petersson, Phys. Rev. Lett. **75**, 4169 (1995) [arXiv:hep-lat/9506025]; Nucl. Phys. B **469**, 419 (1996) [arXiv:hep-lat/9602007].
 - [2] J. Engels, R. Joswig, F. Karsch, E. Laermann, M. Lütgemeier, and B. Petersson, Phys. Lett. B **396**, 210 (1997) [arXiv:hep-lat/9612018]; S. Gottlieb *et al.*, Phys. Rev. D **55**, 6852 (1997) [arXiv:hep-lat/9612020]; C. W. Bernard *et al.* [MILC Collaboration], Phys. Rev. D **55**, 6861 (1997) [arXiv:hep-lat/9612025].
 - [3] F. Karsch, E. Laermann and A. Peikert, Phys. Lett. B **478**, 447 (2000) [arXiv:hep-lat/0002003].
 - [4] A. Peshier, B. Kämpfer, O. P. Pavlenko, and G. Soff, Phys. Rev. D **54**, 2399 (1996); P. Levai and U. W. Heinz, Phys. Rev. C **57**, 1879 (1998) [arXiv:hep-ph/9710463].
 - [5] J. Frenkel, A. V. Saa, and J. C. Taylor, Phys. Rev. D **46**, 3670 (1992); R. Parwani and H. Singh, Phys. Rev. D **51**, 4518 (1995) [arXiv:hep-th/9411065].
 - [6] E. Braaten and A. Nieto, Phys. Rev. D **51**, 6990 (1995) [arXiv:hep-ph/9501375].
 - [7] P. Arnold and C. x. Zhai, Phys. Rev. D **50**, 7603 (1994) [arXiv:hep-ph/9408276]; Phys. Rev. D **51** (1995) 1906 [arXiv:hep-ph/9410360]; C. x. Zhai and B. Kastening, Phys. Rev. D **52**, 7232 (1995) [arXiv:hep-ph/9507380].
 - [8] E. Braaten and A. Nieto, Phys. Rev. Lett. **76**, 1417 (1996) [hep-ph/9508406]; E. Braaten and A. Nieto, Phys. Rev. D **53** (1996) 3421 [hep-ph/9510408].
 - [9] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. D **61**, 014017 (2000) [arXiv:hep-ph/9905337]; J. O. Andersen, E. Braaten, E. Petitgirard and M. Strickland, arXiv:hep-ph/0205085.
 - [10] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. D **63**, 105008 (2001) [arXiv:hep-ph/0007159].
 - [11] J. I. Kapusta, Finite-temperature field theory (Cambridge University Press, Cambridge, UK, 1989); M. Le Bellac, Thermal field theory (Cambridge University Press, Cambridge, UK, 1996).
 - [12] F. Karsch, A. Patkos, and P. Petreczky, Phys. Lett. B **401**, 69 (1997) [arXiv:hep-ph/9702376].
 - [13] S. Chiku and T. Hatsuda, Phys. Rev. D **58**, 076001 (1998) [arXiv:hep-ph/9803226].
 - [14] J. P. Blaizot, E. Iancu and A. Rebhan, Phys. Lett. B **470**, 181 (1999) [arXiv:hep-ph/9910309]; Phys. Rev. Lett. **83**, 2906 (1999) [arXiv:hep-ph/9906340]; Phys. Rev. D **63**, 065003 (2001) [arXiv:hep-ph/0005003]; Phys. Lett. B **523**, 143 (2001) [arXiv:hep-ph/0110369].
 - [15] A. Peshier, Phys. Rev. D **63**, 105004 (2001) [arXiv:hep-ph/0011250].
 - [16] I. T. Drummond, R. R. Horgan, P. V. Landshoff and A. Rebhan, Nucl. Phys. B **524**, 579 (1998) [arXiv:hep-ph/9708426].
 - [17] I. T. Drummond, R. R. Horgan, P. V. Landshoff and A. Rebhan, Phys. Lett. B **460**, 197 (1999) [arXiv:hep-th/9905207].
 - [18] George A. Baker, Jr. and Peter Graves-Morris, *Padé Approximants*, 2nd edition, (Encyclopedia of Mathematics and Its Applications, Vol. 59), edited by Gian-Carlo Rota (Cambridge University Press, 1996).
 - [19] E. Gardi, Phys. Rev. D **56**, 68 (1997) [arXiv:hep-ph/9611453].
 - [20] G. Cvetic, Nucl. Phys. **B517**, 506 (1998); Phys. Rev. D **57**, R3209 (1998); G. Cvetic and R. Kögerler, Nucl. Phys. **B522**, 396 (1998); G. Cvetic, Nucl. Phys. B (Proc. Suppl.) **74**, 333 (1999);
 - [21] G. Cvetic, Phys. Lett. B **486**, 100 (2000); G. Cvetic and R. Kögerler, Phys. Rev. **D63** (2001) 056013.
 - [22] B. Kastening, Phys. Rev. D **56**, 8107 (1997) [arXiv:hep-ph/9708219].
 - [23] T. Hatsuda, Phys. Rev. D **56**, 8111 (1997) [arXiv:hep-ph/9708257].
 - [24] S. J. Brodsky, G. P. Lepage and P. B. Mackenzie, Phys. Rev. D **28**, 228 (1983). A. L. Kataev and V. V. Starshenko, Phys. Rev. D **52**, 402 (1995) [arXiv:hep-ph/9412305]; A. L. Kataev and V. V. Starshenko, Mod. Phys. Lett. A **10**, 235 (1995) [arXiv:hep-ph/9502348]; J. R. Ellis, I. Jack, D. R. Jones, M. Karliner and M. A. Samuel, Phys. Rev. D **57**, 2665 (1998) [arXiv:hep-ph/9710302]. F. A. Chishtie and V. Elias, Phys. Lett. B **521**, 434 (2001) [arXiv:hep-ph/0107052]; C. Contreras, G. Cvetic, K. S. Jeong and T. Lee, arXiv:hep-ph/0203201.
 - [25] D. J. Gross, R. D. Pisarski and L. G. Yaffe, Rev. Mod. Phys. **53**, 43 (1981).
 - [26] E. Braaten, Phys. Rev. Lett. **74**, 2164 (1995) [arXiv:hep-ph/9409434].
 - [27] T. Appelquist and R. D. Pisarski, Phys. Rev. D **23**, 2305 (1981). N. P. Landsman, Nucl. Phys. B **322**, 498 (1989).
 - [28] A. D. Linde, Rept. Prog. Phys. **42**, 389 (1979); A. D. Linde, Phys. Lett. B **96**, 289 (1980).
 - [29] G. Grunberg, Phys. Lett. **95B**, 70 (1980), **110B**, 501(E) (1982); **114B**, 271 (1982); Phys. Rev. D **29**, 2315 (1984).
 - [30] A. L. Kataev, N. V. Krasnikov, and A. A. Pivovarov, Nucl. Phys. **B198**, 508 (1982).
 - [31] A. Dhar and V. Gupta, Phys. Rev. D **29**, 2822 (1984); V. Gupta, D. V. Shirkov, and O. V. Tarasov, Int. J. Mod. Phys. A **6**, 3381 (1991).
 - [32] H. Kleinert, J. Neu, V. Schulte-Frohlinde, K. G. Chetyrkin and S. A. Larin, Phys. Lett. B **272**, 39 (1991) [Erratum-ibid. B **319**, 545 (1993)] [arXiv:hep-th/9503230].




FIG. 1: The Debye screening mass m_E as function of the renormalization scale (RScl) μ_m , when $T=1$ GeV. The upper of the two curves LO TPS and NLO TPS, respectively, has $a(\mu_m^2)$ evolved by the one-loop and two-loop RGE from $a(m_\tau^2)$. All the other curves have $a(\mu_m^2)$ evolved by the four-loop PA [2/3] beta function.




FIG. 2: The F_E (a) and F_M (b) contributions to the free energy F , as functions of the corresponding RScl μ_E , μ_M .




FIG. 3: (a) Various approximants for the normalized sum $R_{E+M} \equiv F_{E+M}/F_{\text{ideal}}$ as functions of the factorization scale Λ_E . (b) shows the separate (E and M) contributions for the stable approximation $R_E[1/1] + R_M[0/2]$. Other details are explained in the text.




FIG. 4: The approximant $R_E[1/1] + R_M[0/2]$ as function of temperature (a) and of $g_s(2\pi T)$ (b), for three different choices of the relevant scales ($\mu_E, \mu_M = \mu_m, \Lambda_E$), as explained in the text. For comparison, the TPS curve in powers of $g_s(2\pi T)$ is included. In our case we have $g_s(2\pi T) \approx 1.2$ (1.0) for $T = 10$ (100) GeV.




FIG. 5: Same as in Fig. 4, but for five different choices of the number of active massless quarks: $n_f = 0, 2, 3, 4$, and 6. The choice of the relevant scales is mid Λ , as explained in the text. In all the previous figures, we took the canonical value $n_f = 3$.




FIG. 6: The screening mass m_L in the massless scalar ϕ^4 theory as a function of the RScI μ_m , at $T=1$ GeV. Other details are given in the text.




FIG. 7: The high energy F_S (a) and low energy F_L (b) contributions to the free energy in ϕ^4 theory as functions of the corresponding RScI's. Other details are given in the text.




FIG. 8: Various approximants for the normalized sum $R_{S+L} \equiv F_{S+L}/F_{\text{ideal}}$ as functions of the factorization scale Λ_F . Other data given in the figure and in the text.




FIG. 9: The normalized sum $R_{S+L} \equiv F_{S+L}/F_{\text{ideal}}$ as a function of $g(2\pi T)$ (a). (b) presents variation of the resummed result $R_S[1/1] + R_L[0/2]$ for three choices of the relevant scales ($\mu_S, \mu_L = \mu_m, \Lambda_F$), as explained in the text.